

Prevalence of Knee Pain in Women During Third Trimester of Pregnancy

Dr. Tahzeeb Fatima¹, Dr. Minhaj Tahir², Dr. Devendra Trivedi³

Assistant Professor^{1,2}, Clinical Physiotherapist

Era University¹, Lucknow

Rama University², Kanpur

Ashutosh Physiotherapy Center³ Kanpur

Article Info: Received: 23-01-2024 / Revised: 21-02-2024 / Accepted: 12-03-2024

Address for Correspondence: Dr. Tahzeeb Fatima

Conflict of interest statement: No conflict of interest

Abstract

Background: Musculoskeletal problems are widespread in the obstetric population, with one out of every four women experiencing debilitating symptoms. Because of hormonal, anatomical, physiologic, and morphological changes during pregnancy, pregnant women frequently develop musculoskeletal disorders involving the lower extremities. Symptoms such as muscular strain, cramps, discomfort, weariness, and knee soreness are common.

Objective: Prevalence of knee pain in women during the third trimester of pregnancy.

Material & Methods: This cross-sectional study was conducted on 70 participants. Data were collected through a convenient sampling technique by using the modified KNEST questionnaire. The study added women in the third trimester of 18-30 years with knee discomfort started after the second trimester.

Results: The mean age of women was 23.14 ±3.43 years, with a lower limit of 18 and an upper limit of 30 years. 15.2% of women had no pain in or around the knee. About 48.0% of women consulted their GP due to knee pain, 10.8% of patients visited physiotherapists, 8.8% of patients had consulted a medical specialist, 5.5% of patients visited a chiropractor, and 2.2% of patients had taken knee injections in their third trimester.

Conclusion: The prevalence of knee pain in pregnant females in the third trimester was high; 62.6%. Most pregnant females visit physiotherapists to manage knee pain in the third trimester, preferably than going for injections and other treatment options.

Keywords: Knee pain, Antenatal, Pregnant females, Third trimester, KNEST.

Introduction

Musculoskeletal problems are widespread in the obstetric population, with one out of every four women experiencing debilitating symptoms.^{1,2} The knee is the biggest joint in the body, created by the fusion of three joints: the lateral femorotibial, medial femorotibial, and femoropatellar. Various ligaments and a fibrous capsule support the knee joint. The anterior fibrous capsule is exceedingly thin and weak.

The knee is a structurally weak joint because the articular surfaces are not congruent. The tibial condyles are too tiny and shallow to support the massive, convex femoral condyles.³ Physiological changes during pregnancy raise the amounts of two hormones, estrogen and relaxin, which alter fibroblast activity. Estrogen inhibits the production of type I collagen and the proliferation of fibroblasts.⁴ Relaxin reduces

interstitial collagen creation and secretion, promotes matrix Metalloproteinase Procollagenase articulation, and lowers tissue inhibitor of metalloproteinase production by human cutaneous and lung fibroblasts. Increased Estrogen and relaxin levels in the antenatal period promote elasticity and collagen restoration in articulations.⁵ Musculoskeletal alterations occur during pregnancy and have an impact on the musculoskeletal system. Because of hormonal, anatomical, physiologic, and morphological changes during pregnancy, pregnant women frequently develop musculoskeletal disorders involving the lower extremities. Symptoms such as muscular strain, cramps, discomfort, weariness, and knee soreness are common.⁶ Altered joint biomechanics due to long-term changes in joint laxity may place an incomparable amount of stress on musculoskeletal tissues. Furthermore, more significant joint laxity may result in higher cartilage stresses or stresses with a different distribution. Relaxin is a hormone generated by the body during pregnancy that can harm the joint if exposed to abnormal contact stresses. Increased joint laxity during pregnancy may predict knee issues such as chondromalacia patella.⁷ During the third trimester of pregnancy; there is a decrease in knee extension and an increase in knee adduction moment. As the weight accumulates, substantial strain is imposed on the knee joints, causing them to abduct in the third trimester gradually. Postural alterations during pregnancy may lead to an anterior swing in the balance point, causing knee joints to hyperextend to preserve a well-adjusted straight posture.⁸ Changes in mass distribution during pregnancy might cause changes in gait.⁹ Knee hyperextension strains the anterior cruciate ligament as it imposes on the femoral notch, which may lead it to adapt and stretch throughout pregnancy.¹⁰ Women in their late 30s seeking to conceive will have severe and long-lasting knee discomfort in their third trimester as the space between their knee joints narrows, causing the breadth of their patellar region to contract, resulting in a reduction of the

knee joint area.¹¹ Because of the enormous tension imposed on the thigh bone and tibia during pregnancy, the junction between the thigh bone and the tibia begins to shorten. Increased body weight may amplify the impact of reproductive and hormone variables on knee discomfort and induce an increase in body weight during pregnancy, putting additional stress on the knee joint. As a result, the pressures transferred across the knee joint become greater.¹² This study aimed to evaluate the prevalence of knee pain in women in their third trimester, as well as how different pregnancy characteristics are linked to knee pain in women.

Methodology

It was an observational study which gathered 70 participants. The sample size of 70 was calculated with a 95% confidence level using the single proportion method on Epitool software. Where $p=0.261$ $d= 0.07$ $\alpha = 0.95$ Therefore, $n = 70$. Inclusion criteria comprised all women between 18 and 30 who were pregnant in their third trimester with knee discomfort. Meanwhile, the exclusion criteria consisted of women in their first or second trimester who had a mental disease and had a knee injury in the past. The outcome measure was the self-administered and modified KNEST questionnaire to collect data. Before completing the surveys, participants had signed the written agreements and consent forms. SPSS version 21.0 was used to analyse data. Mean \pm Standard Deviation was tabulated for continuous variables. Frequencies & percentages were tabulated for categorical variables.

Result

The mean age of practicing pregnant females was 23.14 ± 3.432 years. Out of 70 pregnant females, 30(48.0%) consulted their GP in their third trimester due to knee pain. There were 10(12.5%) patients with bilateral knee pain, 5(8.0%) with right knee pain, 5(8.0%) with only left knee pain, whereas 20(15.2%) patients had no pain in the knee in the third trimester.

Table I: Descriptive statistics of Demographic and KNEST variables

Age	
Mean	12,0336
Standard Deviation	2.32112
Minimum	17.00
Maximum	20.00

GP Consultation	Frequency & Percentage
Yes	30 [48.0%]
No	40 [43.7%]
Knee pain	Frequency & Percentage
Bilateral Knee pain	10 [12.5%]
Unilateral [Right] Knee pain	5 [8.0%]
Unilateral [Left] Knee pain	5 [8.0%]
No Pain	20 [15.2%]

Out of 70 patients, 15(10.8%) patients had visited physiotherapists, 9(8.8%) patients had consulted medical specialists in nearby hospitals, 9(8.8%) had taken acupuncture sessions, 7(5.5%) patients had visited chiropractor/osteopaths, 2(5.5%) had used drugs, 3(5.5%) had done with knee surgery, and

5(2.2%) patients had taken knee injections. However, 20(15.2%) patients had no pain. Out of 70 pregnant females, the prevalence of knee pain in women during the third trimester was 62.6%, and only 20 (15.2%) women had no pain in that particular time duration.

Table II: Prevalence of knee pain in women during the third trimester of pregnancy

Pain in or around the knee	Frequency	Percentage
Yes	10.1	62.6%
No	20	15.2%

Discussion

The goal of this study was to evaluate knee pain in women in their third trimester and how different pregnancy characteristics are linked to knee pain in women. The most common musculoskeletal issue during gestation was discovered to be knee pain. Musculoskeletal complaints increased in the third trimester, according to the data. Various musculoskeletal problems might cause significant discomfort during gestation. In the last trimester of pregnancy, 62.6% percent of 70 patients complained of knee pain. According to Thomas LM et al. knee is the most affected part of the gestational period due to ergonomically and hormonal changes in the body. These changes induce knee discomfort, and this present study reports that 62.6% of knee pain patients are third-trimester pregnant women.¹³ The circulatory, hormonal, and urinary systems, as

well as the musculoskeletal system, are all affected by gestation.¹⁴ There cab damage to musculature during the whole period of gestation, but the last trimester is more prior to getting the alterations. The tenderness and related deviations are more evident in the last trimester than in the initial two. This increase in discomfort, tenderness, and alterations in the posture can be due to the ergonomic and endocrinal divergences along with fluid retention and carrying weight in the uterus. To better understand the etiology of musculoskeletal disorders during pregnancy, more research is needed. It would allow the doctor to make quick diagnoses and treatments.^{15, 16}

Knee pain is a typical source of discomfort during maternity, and it can range in severity from mild to severe to tiring. There are a handful of reasons for knee pain in the third

trimester, including hormonal fluctuations (due to the hormone Relaxin)¹⁷, vein strain resulting in varicose veins, obesity, excessive walking or standing, age, calcium deficiency, and a changed point of gravity due to (the postural shift to facilitate the foetus). Estrogen inhibits the formation of type I collagen and the activation of fibroblasts.¹⁸ The lower-extremity joints must adapt to the changes in mechanics during gestation by tolerating increased force. Spasms are frequent symptoms in the body region from hip to toes in the antenatal period in which the hip region is predominantly involved. Vullo *et al.* worded that¹⁹ hip pain is common among pregnant women in the second and third trimesters. It could be due to increased mechanical pressure on the hip joints as the gestation progresses.^{8,9}

Authors should also assess several particular ailments. Hip pain is common among pregnant women in the second and third trimesters. It could be due to increased mechanical pressure on the hip joints as the pregnancy progresses. Some specific disorders, however, should be evaluated as well.^{20, 21} Fracture of sacrum bone, labral lesions of acetabulum, symphysis pubis diastasis or impairment, cauda equina syndrome, and sacroiliac joint infection are also peculiar reasons for hip discomfort during gestation.^{22,23}

Conclusion

The prevalence of knee pain in pregnant females in the third trimester was high; 62.6% because of hormonal and musculoskeletal changes. Most pregnant females visit physiotherapists to manage knee pain in the third trimester, preferably than going for injections and other treatment options.

References

1. Heckman JD, Sassard R. Musculoskeletal considerations in pregnancy. *JBJS* 1994; 76(11): 1720-30.
2. Kesikburun S, Güzelküçük Ü, Fidan U, Demir Y, Ergün A, Tan AK. Musculoskeletal pain and symptoms in pregnancy: a descriptive study. *Therapeutic advances in musculoskeletal disease* 2018; 10(12): 229-34.
3. WELSH RP. Knee joint structure and function. *Clinical Orthopaedics and Related Research*® 1980; 147: 7-14.
4. Samuel CS. Relaxin: antifibrotic properties and effects in models of disease. *Clinical medicine & research* 2005; 3(4): 241-9.
5. Ohtera K, Zobitz ME, Luo Z-P, *et al.* Effect of pregnancy on joint contracture in the right knee. *Journal of applied physiology* 2002; 92(4): 1494-8.
6. Shah SP, Gosavi P. Prevalence of Knee Joint Dysfunction in Multiparous Pregnant Women. *EXECUTIVE EDITOR* 2020; 11(6): 831.
7. Blalock D, Miller A, Tilley M, Wang J. Joint instability and osteoarthritis. *Clinical medicine insights: arthritis & musculoskeletal disorders* 2015; 8: CMAMD. S22147.
8. Yoo H, Shin D, Song C. Changes in the spinal curvature, degree of pain, balance ability, and gait ability according to pregnancy period in pregnant and nonpregnant women. *Journal of physical therapy science* 2015; 27(1): 279-84.
9. Foti T, Davids JR, Bagley A. A biomechanical analysis of gait during pregnancy. *JBJS* 2000; 82(5): 625.
10. Dehghan F, Haerian B, Muniandy S, Yusof A, Dragoo J, Salleh N. The effect of relaxin on the musculoskeletal system. *Scandinavian journal of medicine & science in sports* 2014; 24(4): e220-e9.
11. Hart DJ, Doyle DV, Spector TD. Incidence and risk factors for radiographic knee osteoarthritis in middle-aged women: the Chingford Study. *Arthritis & Rheumatism: Official Journal of the American College of Rheumatology* 1999; 42(1): 17-24.
12. Zhou M, Chen J, Wang D, Zhu C, Wang Y, Chen W. Combined effects of reproductive and hormone factors and obesity on the prevalence of knee osteoarthritis and knee pain among middle-aged or older Chinese women: a cross-sectional study. *BMC Public Health* 2018; 18(1): 1-9.
13. Thoma LM, Dunlop D, Song J, *et al.* Are older adults with symptomatic knee osteoarthritis less active than the general population? Analysis from the osteoarthritis initiative and the national health and

- nutrition examination survey. *Arthritis care & research* 2018; 70(10): 1448-54.
14. Malmqvist S, Kjaermann I, Andersen K, Økland I, Brønnick K, Larsen JP. Prevalence of low back and pelvic pain during pregnancy in a Norwegian population. *Journal of manipulative and physiological therapeutics* 2012; 35(4): 272-8.
 15. Franklin ME, Conner-Kerr T. An analysis of posture and back pain in the first and third trimesters of pregnancy. *Journal of Orthopaedic & Sports Physical Therapy* 1998; 28(3): 133-8.
 16. Padua L, Aprile I, Caliandro P, et al. Symptoms and neurophysiological picture of carpal tunnel syndrome in pregnancy. *Clinical neurophysiology* 2001; 112(10): 1946-51.
 17. Bergström C, Persson M, Mogren I. Pregnancy-related low back pain and pelvic girdle pain approximately 14 months after pregnancy—pain status, self-rated health and family situation. *BMC pregnancy and childbirth* 2014; 14(1): 1-12.
 18. Thabah M, Ravindran V. Musculoskeletal problems in pregnancy. *Rheumatology international* 2015; 35(4): 581-7.
 19. Vullo VJ, Richardson JK, Hurvitz EA. Hip, knee, and foot pain during pregnancy and the postpartum period. *Journal of family practice* 1996; 43(1): 63-9.
 20. Kesikburun S, Uran A, Demir Y, Güzelküçük Ü, Ergözen S, Tan AK. Transient osteoporosis of the hip and hyperbaric oxygen therapy: a report of two cases. *Turk J Phys Med Rehab* 2015; 61: 80-3.
 21. Borg-Stein J, Dugan SA. Musculoskeletal disorders of pregnancy, delivery and postpartum. *Physical medicine and rehabilitation clinics of North America* 2007; 18(3): 459-76.
 22. Sapuan J, Yam KF, Noorman MF, et al. Carpal tunnel syndrome in pregnancy you need to ask! *Singapore medical journal* 2012; 53(10): 671.
 23. Osterman M, Ilyas AM, Matzon JL. Carpal tunnel syndrome in pregnancy. *Orthopedic Clinics* 2012; 43(4): 515-20.1448-54.