

Contents lists available at www.ijpba.in

International Journal of Pharmaceutical and Biological Science Archive

Index Copernicus Value: 65.90

Volume 6 Issue 2; March-April; 2018; Page No. 01-05

ETHNOMECINAL USES OF FLEMINGIA STROBILIFERA (L).W.T.AITON

Dr.Shreedevi¹, Vd. B.R.Patel²

¹PhD Scholar, Department of Dravya Guna, Institute for Postgraduate Teaching and Research in Ayurveda, Gujarat Ayurved
University, Jamnagar, Gujarat – 361 008, India.

ABSTRACT

The drug *Flemingia strobilifera*(L).W.Aiton is one among many used medicine in India even though not mentioned in Ayurvedic classical texts. The study is conducted to know the ethnomedicinal uses of *Flemingia strobilifera*(L). Various books on ethno-botany, published research and review articles, journals and webpages were reviewed to collect the information regarding Ethnomedicinal uses of *Flemingia strobilifera*(L). It is found that the drug *Flemingia strobilifera*(L). is being used in 17 different states of India like Gujarat, Assam, Bihar etc. Root, Leaf, Seed, Flower, Whole plant is used in the treatment of different ailments. Almost 24 diseases are treated by this drug. Diseases like epilepsy, hysteria, rheumatism, tuberculosis etc are treated by this drug. Countries like Burma, Malaya, Java, Nepal etc are also using *Flemingia strobilifera*(L). in the treatment of different diseases. Other than medicinal usage, the drug is used as host for lac, ornamental plant, in agroforestry, as firewood, green manure etc.

KEY WORDS: Flemingia strobilifera(L).W.Aiton

INTRODUCTION:

Flemingia strobilifera (L).W.Aiton commonly known as Wild hop and luck plant belongs to the family Fabaceae and subfamily Papilionceae. It is known by the name Kaanphoot in Hindi, Makhiyati in Assamese, Kumalu, Nari baalada honne in Kannada, Kanalam, Kumaanchedi, Kumalu, Ponthorilla, Theri in Malayalam, Bandar, Kaanphooti, Klipti in Marati, Bhatamaase, Bhatavaasi in Nepali and Nalla baddu in Telagu¹. The Generic name Flemingia is assigned in the honour of John Fleming, M.D, F.R.S, Physician General and President of Bengal Medical Board². About 35 species are known to occur mainly in tropics of Asia, 15-18 species of them have been reported from India³. Flemingia strobilifera(L). is an important medicinal plant. In wealth of India it has reported that roots of strobilifera(L). are used by Santal tribe in epilepsy, hysteria, insomnia and to relieve pain⁴.

MATERIALS AND METHODS:

Various books on ethno-botany, published research and review articles, journals and webpages were reviewed to collect the information related to Ethnomedicinal uses of *Flemingia strobilifera*(L).

OBSERVATION AND RESULTS:

It is observed that, various parts of *Flemingia strobilifera(L)*. were used by tribes of India, to treat 24 different disease conditions. Though classical texts didn't mention about the use of this plant for clinical use, in ethno medicinal practice, different parts like leaf, whole plants etc were also used with root. Different parts of *Flemingia strobilifera(L)*. used to treat conditions like epilepsy, hysteria, rheumatism, tuberculosis etc. It is used by various tribes of 17 different states of India i.e. Andrapradesh, Gujarat, Haryana, Madhyapradesh, Assam etc.

² Asst Professor, Dept. of Dravyaguna, Institute for Postgraduate Teaching and Research in Ayurveda, Gujarat Ayurved University, Jamnagar, Gujarat – 361 008. India.

Table 1: Different diseased condition where *Flemingia strobilifera*(L) W.T. Aiton is useful 5-15

Sr. No	Name of disease/ condition	Part used	
1.	Haemorrhage	Entire plant	
2.	Bronchial Asthma Entire plant		
3.	Hysteria	Root, Leaf	
4.	Fever	Root	
5.	Rheumatism	Root	
6.	Worm	Root	
7.	Scabies	Root	
8.	As contraceptive	Seed	
9.	Analgesic	Leaf, Root	
10.	To induce sleep	Root	
11.	As tonic	Root	
12.	Vermifuge	Leaf	
13.	Epilepsy	Root	
14.	Blood Dysentery	Root	
15.	Insect sting Leaf		
16.	Veterinary skin disease	Whole plant	
17.	Gastric complaints	Root	
18.	Leucorrhea	Root	
19.	Oedema Leaf		
20.	Diarrhea Root		
21.	Kidney stone Root		
22.	Fly repellant Root		
23.	Cessation of menstuation born complaints	Leaf	
24.	Tuberculosis	Flower, Leaf	

Table 2: Use of Flemingia strobilifera(L) W.T.Aiton in different diseases in different states of India5-15

Sr. No.	Disease / Disease stage	Part used	Name of the State
1.	As analgesic	Root	Assam
2.	Body ache	Root	Uttar Pradesh, Maharashtra, Madya Pradesh, Andra Pradesh
3.	Epilepsy	Root	Sikkim, Bengal, Bihar, Orissa
4.	Fever	Root	Maharashtra, Madya Pradesh, Andra Pradesh
5.	Gastric Complaints	Root	Sikkim, Bengal, Bihar, Orissa
6.	Hysteria	Root	Sikkim, Bengal, Bihar, Orissa
7.	Insecticidal	Plant part not specified	Maharashtra, Madya Pradesh, Andra Pradesh
8.	Leucorrhea	Root	Maharashtra, Madya Pradesh, Andra Pradesh
9.	Sedative	Root	Assam
10.	Vermifuge	Leaf	Punjab, Haryana, Rajastan, Gujarat, Assam
11.	As Tonic	Leaf	Uttar Pradesh

Method of administration of *Flemingia* strobilifera(L). 16-20

- One tea spoonful of root paste is given twice daily for 3-5 days for the treatment of Epilepsy and Hysteria.
- 3-5 teaspoonful of root paste is given daily for Gastric disorder.
- Root of the plant in the form of paste is applied locally in scabies.
- In haemorrhage decoction of entire plant is administered.
- Even in Bronchial Asthma, decoction is prepared using entire plant and then given orally.
- Lut fresh leaves boiled in coconut oil till the green turns to black, allowed to cool and applied as medicated oil, half an hour before bath for problems associated with body pain.

Medicinal use of *Flemingia strobilifera*(L). in different countries ^{21,22,23}

One seed a month is chewed by women as a contraceptive.

- In India, root paste applied externally to body swelling.
- > Root used for epilepsy in India.
- ➤ In Malaya, decoction of leaves administered after childbirth and for bathing.
- Decoction of leaves used for rheumatism.
- In Java, decoction of leaves used internally and externally as vermifuge for children.
- In Bangladesh leaf juice used to treat worms.
- ➤ In Nepal, Root juice (fresh roots crushed in cup of water by mortar and pestle) is taken twice daily for 7 days for diarrhea and dysentery.
- In Trinidad and Tobago, used for kidney stones.
- ➤ In India leaves and flowers used for tuberculosis. Roots used for hysteria, root juice for diarrhea and dysentery.
- ➤ In India, cut fresh leaves boiled in coconut oil till the green turns to black, allowed to cool and applied as medicated oil half an hour before bath for problems associated with body pain¹⁴.
- ➤ In Burma the roots of F. *Strobilifera*(L). are used to treat epilepsy.

- ➤ The Marma tribe uses this plant as fly repellent.
- Decoction of the leaf is taken orally by Marma tribe to cure body swelling due to cessation of menstruation; bath taken with leaf-boiled water has similar effect. The plant is also used for rheumatic fever.

Other than medicinal uses of *Flemingia* strobilifera(L). ²⁴

In the Philippines, dried bracts used for stuffing pillows and cushions.

Agroforestry

Boundary, barrier or support

Erosion control or dune stabilization

Land reclamation

Revegetation

Soil conservation

Soil improvement

Fuelwood

Green manure

Host for Lac

Mulches

Ornamental

CONCLUSION:

It is observed that, various parts of Flemingia strobilifera(L).W.Aiton. were used by tribes of India, to treat 24 different disease conditions. Though classical texts didn't mention about the use of this plant for clinical use, in ethno medicinal practice, different parts like leaf, seed and whole plants were also used with root. Different parts of Flemingia strobilifera(L). are used to treat conditions like fever, acidity, dysentery, diarrhea, epilepsy, hysteria, dermal disorders etc. It is used by various tribes of 17 different states of India i.e. Andrapradesh, Gujarat, Sikkim, Madhya Pradesh, Assam etc. It is used in different counties in the treatment of diseases. Countries like Java, Burma, Bangla, Nepal etc are using this plant in the treatment of many diseases. Thus the leaf, stem, root etc all the parts are having medicinal value. Since olden time this plant is used for various ailments. Its use as medicine is found throughout India even though not found in classical Ayurvedic texts.

REFERENCE

- 1. Kirtikar KR; Basu BD (1989), Indian Medicinal Plants, Published by Lalit Mohan Basu, Allahabad, India, 2nd Edition. vol. I. p. 813-814.
- **2.** Dr. Anil K. Dhiman , Medicinal Plants of Uttaranchal State, Chowkambha Sanskrit Series, Varanasi, first edition, 2004, pg 179
- **3.** Dr. Anil K. Dhiman , Medicinal Plants of Uttaranchal State, Chowkambha Sanskrit Series, Varanasi, first edition, 2004, pg 179
- **4.** Kirtikar KR; Basu BD (1989), Indian Medicinal Plants, Published by Lalit Mohan Basu, Allahabad, India, 2nd Edition. vol. I. p. 814.
- Dr.S.K.Jain, Dictionary of Indian Folk Medicine and Ethnobotany, Deep Publisher, 1991, p 91,92
- **6.** Baruah, Parkutty & G.C. Sharma, 1987, Studies on the medicinal uses of plants by the Noth-East tribes -3 pg71-76
- **7.** Sharma, B.D & S.K Malhotra, 1984, A contribution to the ethnobotany of tribal areas in Maharashtra, pg 533-537
- **8.** Sharma P.K, S.K Dhyani & V. Shanker 1979, Some useful and medicinal plants of the district Dehradun ans Siwalik, pg 17-43
- **9.** Goel, A.K Sahoo & V. Mudgal, 1984, A contribution of Santal Pargana, Bihar
- **10.** Jain S.K, 1965, Medicinal plantlore of the tribals of Bastar, pg 236-250
- **11.** Jain S.K, Studies in Indian Ethnobotany- 2,1963, Plants used in medicine by the tribals of Madya Pradesh, Bull, pg 126-128
- **12.** Saxena H.O, 1986, Observations on the ethnobotany of Madhya Pradessh, Bull,pg 149-156
- **13.** Singh V & R.P Pandey, 1980, Medicinal plantlore of the tribals of eastern Rajastan, pg 137-147
- **14.** Harja P.K & A. K Baishya, 1981, Ethnobotanical notes on the Miris of Assam, pg 161-169
- **15.** Maheshwari J.K, K.K Singh & S Saha, 1980, Ethnomedicinal uses of plants Tharus in Kheri District, UP, pg 318-337.
- **16.** Maheshwari J.K, Ethnobotany and Medicinal Plants of Indian Subcontinent, Scientific Publishers, Jodhpur, 2003, pg 553

- **17.** Dr. Anil K. Dhiman , Medicinal Plants of Uttaranchal State, Chowkambha Sanskrit Series, Varanasi, first edition, 2004, pg 179
- **18.** A M Saren, R. Sen & D.C Pal, A Contribution to the Ethnobotany of Bankura District, West Bengal, pg 553
- **19.** Mahalakshmi Dutta & Subhan C. Nath, Ethno-Medico-Botany of TAI- Ahoms of Assam, India, pg 596
- **20.** Cross-Cultural Ethnobotanical Studies of Santhal Pargana(Eastern India) and Western Ghats(Southern India), pg 149
- 21. https:/www.cabi.org>isc>datsheet, 22Nov2017
- **22.** https://www.mpbd.info>plants>flemingia-strobilifera
- **23.** http://tropical.theferns.info/viewtropical.php?i d=Flemingia+strobilifera
- 24. https:/www.cabi.org>isc>datsheet, 22Nov2017